



(SUOX)

a c a c c a a b ca

L

L

SUOX

T T

P c a M c a C C

S a a C C


& S a a Sa D B L L C C

B a A b





(OD)





1. 50 5 L 245 L & 50
2. 100 5 L 495 L & 100
3. 1000 5 L 95 L & 20 5 L
20 245 L & 50 1000
4. 100000 5 L 195 L & 40
5 L 40 245 L & 50 5 L 2000
245 L & 50 100000
5. 3 L 100
6. PBS &

O a



L(

) L, C

200

L

L

L

L

L


L

L

L

L

L



L

L

L

L

L

L

L

HRP

L

L

TMB —



C

TMB

TMB

1.

2.

3.



		L			
C				L	
		L			C
			L		
	L			C	
		L			
TMB		L	C		
	L				
				OD	



●

():CV
1

():CV
3

●

SUOX

EDTA		
a		

●

SUOX 2 4 8 16

EDTA				
a				



/

1. 加入 100 μl 待测样本或标准品，轻轻混匀，室温孵育 30 分钟。 2. 加入 100 μl 生物素化抗人 IgG，轻轻混匀，室温孵育 30 分钟。 3. 加入 100 μl 链霉亲和素-生物素复合物，轻轻混匀，室温孵育 30 分钟。 4. 加入 100 μl 链霉亲和素-生物素复合物，轻轻混匀，室温孵育 30 分钟。 5. 加入 100 μl 链霉亲和素-生物素复合物，轻轻混匀，室温孵育 30 分钟。 6. 加入 100 μl 链霉亲和素-生物素复合物，轻轻混匀，室温孵育 30 分钟。 7. 加入 100 μl 链霉亲和素-生物素复合物，轻轻混匀，室温孵育 30 分钟。 8. 加入 100 μl 链霉亲和素-生物素复合物，轻轻混匀，室温孵育 30 分钟。		加入 100 μl 链霉亲和素-生物素复合物，轻轻混匀，室温孵育 30 分钟。
		加入 100 μl 链霉亲和素-生物素复合物，轻轻混匀，室温孵育 30 分钟。
		加入 100 μl 链霉亲和素-生物素复合物，轻轻混匀，室温孵育 30 分钟。
		加入 100 μl 链霉亲和素-生物素复合物，轻轻混匀，室温孵育 30 分钟。
	Avidin-HRP	加入 100 μl 链霉亲和素-生物素复合物，轻轻混匀，室温孵育 30 分钟。
		加入 100 μl 链霉亲和素-生物素复合物，轻轻混匀，室温孵育 30 分钟。
		加入 100 μl 链霉亲和素-生物素复合物，轻轻混匀，室温孵育 30 分钟。
		TMB 450nm 650nm





		30min
		15-30min
		PH
		(
	NaN ₃	NaN ₃
		TMB 450nm 650nm



/

		,
		,
		10
		,
		,
		,
		,
		200 µL/
		,
		,
	,	,
		,
		,
		3 , 2 (2)



		3 , ,
		, , , ,
		, , , ,
	, ,	, , , ,
	, ,	6min, 3000 rpm
	, ,	, , , ,



ELK B c

027-59760950

elkbio@elkbiotech.cn

